

In vitro free radical scavenging effect of *Butea monosperma* leaf extracts

Bappa Sarkar¹,
Payel Manna¹,
Sanjib Bhattacharya^{2*},
Moulisha Biswas¹

¹Bengal Institute of
Pharmaceutical Sciences, Kalyani,
Nadia 741235, West Bengal, India

²Faculty of Pharmaceutical
Sciences, PCTE Group of Institutes,
Ludhiana 142021, Punjab, India

J. Adv. Pharm. Edu. & Res.

ABSTRACT

Butea monosperma commonly known as *Palash* in Sanskrit and Bengali and Flame of the Forest in English is a medium-sized dry season deciduous tree indigenous to tropical and sub-tropical parts of the Indian Subcontinent and Southeast Asia. The present investigation assessed the different solvent extracts of *B. monosperma* leaf for their *in vitro* free radical scavenging potential by 1,1-diphenyl-2-picryl-hydrazyl (DPPH) radical scavenging assay. All of the extracts exhibited potent *in vitro* free radical scavenging activity that increased with extract concentrations. The methanol extract was found to be the most potent in this regard, followed by the chloroform and petroleum ether extracts respectively. Therefore, the present study confirms marked *in vitro* free radical scavenging activity of *Butea monosperma* leaf.

Keywords: Free radical scavenging, leaf, antioxidant, *Butea monosperma*.

INTRODUCTION

Antioxidants protect living organisms from damage caused by uncontrolled production of reactive oxygen species (ROS) and the concomitant lipid peroxidation, protein damage and DNA strand breaking. Current interest is focused on the potential role of antioxidants and antioxidant enzymes in the treatment and prevention of atherosclerosis, heart failure, neurodegenerative disorders, aging, cancer, diabetes mellitus and several others diseases. [1]

Antioxidants are supplemented to a variety of foods to prevent free radical induced lipid peroxidation, which is responsible for the development of off-flavors and the undesirable chemical compounds in food. [2] These ROS cause destructive and irreversible damage to the components of a cell, such as lipids, proteins DNA and other macromolecules. Although normal cells possess antioxidant defense systems against ROS in the cells induces diseases such as cancer and aging. [3]

ROS are formed and degraded by all aerobic organisms. ROS can readily react with most

biomolecules including proteins, lipids, lipoproteins and DNA. Exogenous chemical and endogenous metabolic processes in the human body or in the food system might produce highly reactive oxygen species, which are capable of oxidizing biological molecules, resulting in tissue damage and cell death. When the mechanism of antioxidant protection becomes unbalanced by exogenous and endogenous factors, it results in inflammation, diabetes, genotoxicity, cancer and accelerating aging. [4]

Antioxidant supplements or foods containing antioxidants may be used to help the human body reduce oxidative damage. The most commonly used antioxidants are BHA, BHT, propyl gallate and *tert*-butyl-hydroquinone. [5] However, they have been suspected of being responsible for liver damage and carcinogenesis in laboratory animals. Therefore, the development and use of more effective antioxidants is desired.

Butea monosperma (Lam.) Taub. (Family: Leguminosae: Papilionoideae), commonly known as *Palash* in Sanskrit and Bengali and Flame of the Forest in English is a medium-sized dry season deciduous tree indigenous to tropical and sub-tropical parts of the Indian Subcontinent and Southeast Asia. It is grown throughout the Indian subcontinent as ornamental plant for its bright orange flowers and

Address for correspondence

Dr. Sanjib Bhattacharya
Faculty of Pharmaceutical Sciences,
PCTE Group of Institutes, Ludhiana 142021, Punjab, India.
E-mail: sakkwai@yahoo.com

Access this article online
www.japer.in

timber. Traditionally, all different part of this plant, namely leaf, root, fruit, bark and flower are used in treatment of various kinds of disorders in humans. [6] However, reports on the experimental studies on this plant especially on its leaf part are comparatively scanty. In the present investigation, we have aimed to evaluate *in vitro* free radical scavenging activity of different solvent extracts from *B. monosperma* leaf against 1, 1-diphenyl-2-picryl-hydrazil radicals to assess the possible antioxidant effect.

MATERIALS AND METHODS

Plant material

The mature leaves of *Butea monosperma* (Lam.) Taub. (Family: Leguminosae: Papilionoideae), were collected during the month of August 2014 from Kalyani, Nadia, West Bengal, India. The plant species was taxonomically identified at the Central National Herbarium, Botanical Survey of India, Howrah, West Bengal, India. The voucher specimen [CNH/Tech. II/2014/72/150] was maintained in our research laboratory for future reference. Just after collection the fresh leaves were cleaned thoroughly with running tap water and shade-dried at room temperature (24-26°C) with occasional shifting and then powdered with mechanical grinder, passing through sieve no. 40, and stored in an air-tight container for use in the study.

Preparation of plant extracts

The dried powdered material was successively extracted with petroleum ether, chloroform and methanol for 72 h at room temperature (24-26°C) in a cone shaped percolator. Then the solvent was distilled off in reduced pressure and resulting semisolid mass was vacuum dried using rotary vacuum evaporator to yield the solid extracts and stored in refrigerator for use in the present study. Preliminary phytochemical studies were performed on the three extracts. [7]

Chemicals

L ascorbic acid (vitamin C) and 1,1-diphenyl-2-picryl-hydrazyl (DPPH) from Sigma Chemical Co. Ltd. (St. Louis, MO, USA). All the other chemicals and reagents

were of analytical grade obtained commercially from Merck.

Free radical scavenging activity assessed by 1, 1-diphenyl-2-picryl-hydrazil

The free radical scavenging activity of all of three extracts were measured by 1,1-diphenyl-2-picryl-hydrazil (DPPH) using the reported method. [8] Briefly, an 0.1 mM solution of DPPH in methanol was prepared, and 1 ml of this solution was added to 3 ml of the all of the extracts solution respectively in chloroform, petroleum ether and methanol at different concentrations (2, 4, 6, 8, 10, 15 µg/ml). The mixture were shaken vigorously and allowed to stand at room temperature for 30 min. Then the absorbance was measured at 517 nm using a UV-Vis spectrophotometer (Genesys 10 UV: Thermo Electron Corporation). Lower absorbance values of reaction mixture indicate higher free radical scavenging activity. Ascorbic acid at same concentrations was used as reference agent. The capability to scavenge the DPPH radical was calculated by using the following equation:

$$\text{DPPH scavenging effect (\%)} = [(A_0 - A_1) / A_0] \times 100$$

Where, A_0 is the absorbance of the control reaction, and A_1 is the absorbance of presence of all of the extract samples or reference. The results are stated in Table 1.

RESULTS AND DISCUSSION

Traditional medicine worldwide is being reevaluated by extensive research on different plant species and their therapeutic principles. Plants produce antioxidants to control the oxidative stress caused by chemicals, sunbeams and oxygen species; they can represent a source of new compounds with antioxidant activity. [9-11]

In the present study, the preliminary phytochemical studies indicated the presence of triterpenoids and steroids in petroleum ether extract; triterpenoids, steroids, glycosides and alkaloids in chloroform extract; carbohydrates and polyphenols in methanol extract.

The stable DPPH radical model is a widely used, relatively quick and precise method for the evaluation of free radical scavenging activity. The effects of antioxidants on DPPH radical scavenging is thought to be due to their hydrogen-donating ability. DPPH is a stable free radical that accepts an electron or hydrogen radical to become a stable diamagnetic molecule. The absorption maximum of a stable DPPH radical in methanol was 517 nm. The decrease in absorbance of DPPH radical caused by antioxidants, because of the reaction between antioxidant molecules and radical progressed, results in the scavenging of the radicals by hydrogen donation. It is visually noticeable that as a change in color from purple to yellow. Hence, DPPH is usually used as a substrate to evaluate the antioxidant activity of different antioxidants. [12, 13] It has been reported that oxidative stress, which occurs when free radical formation exceeds the body's ability to protect itself, forms the biological basis of several chronic disease conditions such as arteriosclerosis, cancer etc. [14]

Based on the data obtained from the present work, all the test extracts were effective free radical inhibitor or scavenger, which reacts with free radicals, which may limit free radical damage occurring in the human body. The results are summarized in the Table 1. Free radical scavenging activity *in vitro* also increased with increasing concentration in the range of 2-15 µg/ml indicating concentration dependence on activity. The methanol extract was found to demonstrate the most active free radical scavenging potential at the test concentrations, followed by the chloroform and petroleum ether extracts respectively. The presence of polyphenolic compounds only in the methanol extract may be responsible for its maximum antioxidative potential, as polyphenols are well known as natural antioxidants. [15, 16] The present preliminary study, therefore, confirms remarkable *in vitro* free radical scavenging activity of *Butea monosperma* leaf against DPPH.

Table 1: DPPH scavenging effect of the different extracts of *B. monosperma* leaf.

Extracts	Concentrations (µg/ml)	% of DPPH scavenging activity respectively	Mean ± SEM
Petroleum ether	2, 4, 6, 8, 10, 15	27.63, 29.18, 33.27, 54.63, 67.45, 78.83	48.49 ± 4.27
Chloroform	2, 4, 6, 8, 10, 15	31.42, 38.59, 51.18, 66.51, 72.61, 85.44	57.63 ± 4.63
Methanol	2, 4, 6, 8, 10, 15	42.75, 58.27, 77.24, 81.92, 87.27, 91.45	73.15 ± 5.45
Ascorbic acid	2, 4, 6, 8, 10, 15	25.70, 32.10, 42.50, 87.54, 92.20, 97.23	62.87 ± 5.29

ACKNOWLEDGEMENT

The authors are thankful to the authority of Bengal Institute of Pharmaceutical Sciences Kalyani, Nadia 741235, West Bengal, India for necessary facilities for the present study.

REFERENCES

- Ajitha M., Rajnarayana K. Role of oxygen free radicals in human disease. *Indian Drugs*. 2001; 38: 545-554.
- Halliwell B., Aeschbach R., Loliger J., Aruom O.I. The characterization of antioxidants. *Food Chem. Toxicol*. 1995; 33: 601-617.
- Mates J.M., Sanchez-Jimenez F.M. Role of reactive oxygen species in apoptosis: Implications for cancer therapy. *Int. J. Biochem. Cell. Biol*. 2000; 32: 57-170.
- Buyukokuroglu M.E., Gulein L., Oktav M., Kufrevioglu O.I. *In vitro* antioxidant properties of dantrolene sodium. *Pharmacol. Res*. 2001; 44: 491-495.
- Gulcin I., Beydemir S., Ahmet A.H., Mahfuz E., Emin B.M. *In vitro* antioxidant properties of morphine. *Pharmacol. Res*. 2004; 49: 59-66.
- Yadav S., Patgiri B.J., Prajapati P.K. Review of Bio-active Principle of *Butea monosperma* (Lam.) Kuntze. *Indian J. Ethno-Phytopharm*. 2015; 1: 45-51.
- Kokate C.K. *Practical Pharmacognosy*. 4th Edition. New Delhi. Vallabh Prakashan. 1994; 107-112.
- Blois M.S. Antioxidant determinations by the use of stable free radical. *Nature*. 1958; 26: 1199-1200.
- Bhattacharya S., Nagaich U. Assessment of anti-nociceptive efficacy of *Costus speciosus* rhizome in

- Swiss albino mice. *J. Adv. Pharm. Technol. Res.* 2010; 1: 34-40.
10. Bhattacharya S., Roy B. Preliminary investigation of antipyretic activity of *Cuscuta reflexa* in rats. *J. Adv. Pharm. Technol. Res.* 2010; 1: 83-87.
11. Dey P., Chandra S., Chatterjee P., Bhattacharya S. Neuropharmacological properties of *Mikania scandens* (L.) Willd. (Asteraceae). *J. Adv. Pharm. Technol. Res.* 2011; 2: 255-259.
12. Bhattacharya S. Natural antimutagens: A review. *Res. J. Med. Plants.* 2011; 5: 116-126.
13. Bhattacharya S., Mueen Ahmed K.K., Chakraborty S. Free radicals and cardiovascular diseases: An update. *Free Rad. Antiox.* 2011; 1: 17-22.
14. Bhattacharya S. Anticarcinogenic property of medicinal plants: involvement of antioxidant role. In: Capasso A. Ed. *Medicinal Plants as Antioxidant Agents: Understanding their Mechanism of Action and Therapeutic Efficacy.* Trivandrum, India: Research Signpost. 2012; 83-96.
15. Bhattacharya S. Are we in the polyphenols era? *Pharmacogn. Res.* 2011; 3: 147.
16. Chatterjee P., Chandra S., Dey P., Bhattacharya S. Evaluation of anti-inflammatory effects of green tea and black tea: A comparative *in vitro* study. *J. Adv. Pharm. Technol. Res.* 2012; 3: 136-138.

How to cite this article: Bappa Sarkar¹, Payel Manna¹, Sanjib Bhattacharya^{2*}, Moulisha Biswas¹; *In vitro* free radical scavenging effect of *Butea monosperma* leaf extracts; *J. Adv. Pharm. Edu. & Res.* 2015; 5(3): 103-106.

Source of Support: Nil, **Conflict of Interest:** Nil